

Prevalence and Genotype Distribution of HPV in Cytology Specimens Containing Atypical Glandular Cells: A Case Control Study

S. B. Namugenyi, M. J. Balsan, S. N. Glick, J. A. Jordan

The George Washington University School of Public Health and Health Sciences, Washington DC

School of Public Health
& Health Services

THE GEORGE WASHINGTON UNIVERSITY

Introduction

- HR-HPV responsible for $\geq 99\%$ of all cervical cancer, with HPV 16/18 causing $\sim 70\%$
- Pap test: Marked decline in the incidence of squamous cell cervical cancers, but incidence of glandular cell cancers has increased^{1, 2}
- Atypical glandular cell (AGC) grade Pap test results are rare (0.2-0.7%) but 38% represent a corresponding high-grade lesion on biopsy^{3, 4}
- AGC management guidelines: Colposcopy, endocervical sampling & HR-HPV DNA testing⁵

Objective

- Determine prevalence & genotype distribution of HPV in AGC-grade cytology specimens compared to women whose Pap smears were negative for intraepithelial lesion or malignancy (NILM), with ages known for both groups of women.

Methods

- Collected residual, de-identified AGC & NILM cytology specimens
- DNA was extracted using QIAamp MinElute Media Kit and analyzed by PCR using Linear Array HPV Genotyping and Detection Test Kits
- Multivariate logistic regression compared HPV prevalence and genotype distribution between cases and controls to generate age-adjusted odds ratios (ORadj) and 95% confidence intervals (CI)

Results

Table 1. Study Sample Description

	Total number tested	Mean age (yr)	Median age (yr)	Age range (yr)
AGC-grade cases	53	57	56	18-95
NILM-grade controls	338	45	43	20-91

Table 2 A. Percent of specimens containing detectable levels of HR-HPV DNA

	AGC-grade cases	NILM-grade controls
HR-HPV Status *	%	%
Positive	34	7.4
Negative	66	92.6
Total	100	100

*Regardless of LR- HPV co-infection

Table 2 B. Percent of specimens containing detectable levels of LR-HPV DNA

	AGC-grade cases	NILM-grade controls
LR HPV Status *	%	%
Positive	13.2	17.2
Negative	86.8	82.8
Total	100	100



*Regardless of HR-HPV co-infection

Table 3. Multivariate age-adjusted logistic regression analysis comparing risk of finding HR-HPV or LR-HPV DNA in AGC-grade cases compared to NILM-grade controls

HPV Type(s)	*ORadj (95% CI)	P value
Any HR HPV (inc. 16/18)	9.11 (4.08, 20.33)	< 0.001
HR HPV 16/18 only [‡]	40.10 (10.73, 149.88)	< 0.001
Any LR HPV	0.91 (0.35, 2.31)	0.834

* ORadj; adjusted Odds Ratio, 95% CI; 95% Confidence Interval,

[‡]; Irrespective of other HR HPV or LR HPV types



Conclusion

- AGC-grade cases contained a significantly higher rate of HR-HPV, especially HPV types 16/18 when compared to NILM controls
- Prevalence of LR-HPV: Not significantly different between cases and controls – suggesting sexual behavior practices similar between both groups e.g. unprotected intercourse
- Most cases (78%) and controls (74%) had single infections
- When multiple HPV types: More common in AGC cases (5.7%) than NILM controls (0.6%)
- Findings support guidelines that HPV testing should be performed on specimens with AGC-grade diagnosis
- HPV 16/18 genotyping may be valuable for managing women with AGC-grade Pap test results

References

1. Lai CR, et al. (2012) Available from: <http://www.intechopen.com/books/intraepithelial-neoplasia/the-role-of-the-pap-smear-diagnosis-atypical-glandular-cells-agc>. pp. 365-380.
2. Smith HO, et al. (2000) Gynecol Oncol 78(2): 97-105.
3. Chen SF, et al. (2005) Gynecol Oncol 99: 578-584.
4. Apgar BS, et al (2009) Am Fam Physician 80(2): 147-155.
5. Wright TC Jr, et al. (2007) Am J Obstet Gynecol 197(4): 346-355.

Funding

Funding provided by Roche Diagnostics

HR-HPV types: 16, 18, 26, 31, 33, 35, 39, 45, 51, 52, 53, 56, 58, 59, 66, 68, 73, 82
LR-HPV types: 6, 11, 40, 42, 54, 55, 61, 62, 64, 67, 69, 70, 71, 72, 81, 83, 84, IS39, CP6108